

On Solvability of Sylvester Equation via Coupled Fixed Point

Lourabi Hariz Bekkar, Abdelouahab Mansour

¹*Mathematics Department, Faculty of Exact Sciences, El-Oued University,
P.O.Box789, El-Oued 39000, Algeria
harizlnd@yahoo.fr, mansourabdelouahab@yahoo.fr*

Keywords: Sylvester equation, Hermetian solution, coupled fixed point.

In this paper, we will study existence and uniqueness of solution to Sylvester equation

$$AX - XB = C$$

where C is hermetian positive definite matrix and A, B an arbitrary matrices, by using coupled fixed point technique.

References

- [1] R. H. Bartels, G. W. Stewart, Solution of the equation $AX + XB = C$, Comm. ACM. 15 (9) (1972) 820–826.
- [2] T. G. Bhaskar, V. Lakshmikantham, Fixed point theory in partially ordered metric spaces and applications, Nonlinear Anal. 65 (2006) 1379–1393.
- [3] A. Mansour, L. Hariz and H. Gaaya, *A priori estimate for the solution of sylvester equation*, J. Adv. Maths, 10(7) (2015), 3633–3638.
- [4] S. Mecheri and A. Mansour, On the operator equation $AXB - XD = E$, Lobachevskii journal of mathematics 30 (NTA3), 224 (2009).
- [5] A. C. M. Ran, M. C. B. Reurings, *A fixed point theorem in partially ordered sets and some applications to matrix equations*, Proc. Amer. Math. Soc. 132 (2004) 1435–1443.