

Solvers of Optimal Complexity: from Sparse SPD Systems to a Class of Non-Local Problems

Svetozar Margenov

*Institute of Information and Communication Technologies
Bulgarian Academy of Sciences
Acad. G. Bonchev St, Bl. 25A, 1113 Sofia, Bulgaria
margenov@parallel.bas.bg*

Keywords: Finite Element Methods, Numerical Linear Algebra, Multilevel Solvers, Fractional Powers of SPD Matrices

Hierarchical basis algebraic multilevel iteration (AMLI) methods for elliptic problems are introduced at the beginning, assuming that the coefficient jumps are aligned with the coarsest triangulation. The obtained condition number estimates are uniform with respect to mesh and coefficient anisotropy, coefficient jumps, and number of degrees of freedom [1]. More recently, robust multilevel methods for high-frequency and high-contrast problems are developed. Some advantages of the nonlinear AMLI are discussed in this context. The auxiliary space multigrid (ASMG) method for highly heterogeneous media, based on preconditioning of the related weighted $H(\text{div})$ -norm is presented at the end [2]. The second part of the talk is motivated by the active on-going research in fractional diffusion problems. Several different techniques are proposed to localize the non-local operator, thus increasing the space dimension of the original computational domain. An alternative approach is developed in [3]. The main goal is to reduce the computational complexity. Let A be a properly scaled symmetric and positive definite (SPD) matrix. A method for solving algebraic systems of linear equations involving A^α , $0 < \alpha < 1$ is discussed. The solver is based on best uniform rational approximations of the scalar functions. Although the fractional power of A is a dense matrix, the algorithm has complexity of order $O(N)$, where N is the number of unknowns. Some parallel scalability results are provided at the end.

References

- [1] Kraus, J., Margenov, S., Robust Algebraic Multilevel Methods and Algorithms, De Gruyter Incorporated, Walter, Berlin-New York, 2009
- [2] Kraus, J., Lazarov, R., Lymbery, M., Margenov, S., Zikatanov, L., Preconditioning Heterogeneous $h(\text{div})$ Problems by Additive Schur Complement Approximation and Applications, SIAM Journal on Scientific Computing, 38(2), A875–A898, 2016
- [3] Harizanov, S., Lazarov, R., Margenov, S., Marinov, P., Vutov, Y., Optimal solvers for linear systems with fractional powers of sparse SPD matrices (submitted), available as arXiv:1612.04846v1, 2016